

# Perfect 1-Factorisations



David A. Pike  
Memorial University of Newfoundland

## Definition:


A  $k$ -factor of a graph  $G$  is a  $k$ -regular spanning subgraph of  $G$ .

A  $k$ -factorisation of a graph  $G$  is a partition of the edge set  $E(G)$  of  $G$  into  $k$ -factors.


Observe: if  $G$  admits a  $k$ -factorisation then  $G$  is regular and  $k$  must divide the degree  $\Delta$ .

Also, any graph that has a 1-factorisation must have an even number of vertices.


# Example: a 1-Factorisation of $K_8$




$F_0$




$F_1$




$F_2$




$F_3$



$F_4$



$F_5$



$F_6$

## 1-Factorisation Conjecture (Dirac, 1950s?)

Suppose  $G$  is a regular graph of even order.

If  $\Delta \geq \frac{1}{2}|V(G)|$  then  $G$  has a 1-factorisation.

## Theorem (Cariolaro and Hilton, 2009)

Suppose  $G$  is a regular graph of even order.


If  $\Delta \geq \frac{1}{6}(\sqrt{57} - 3)|V(G)|$  then  $G$  has a 1-factorisation.

## Theorem (Walecki, 1890s)

For each  $m \geq 2$ , the complete graph  $K_{2m}$  has a 1-factorisation.


# Theorem (Walecki, 1890s)

For each  $m \geq 2$ , the complete graph  $K_{2m}$  has a 1-factorisation.




# Theorem (Walecki, 1890s)

For each  $m \geq 2$ , the complete graph  $K_{2m}$  has a 1-factorisation.



# Theorem (Walecki, 1890s)


For each  $m \geq 2$ , the complete graph  $K_{2m}$  has a 1-factorisation.



For each  $i = 1, 2, \dots, 2m - 2$ ,  
let  $F_i = F_0 + i \pmod{2m - 1}$ .

# Theorem (Walecki, 1890s)

For each  $m \geq 2$ , the complete graph  $K_{2m}$  has a 1-factorisation.




$F_0$

This 1-factorisation  
is denoted  $GK_{2m}$


For each  $i = 1, 2, \dots, 2m - 2$ ,  
let  $F_i = F_0 + i \pmod{2m - 1}$ .




# The 1-Factorisation of $K_8$ known as $GK_8$




$F_0$




$F_1$




$F_2$




$F_3$



$F_4$



$F_5$



$F_6$

## Definition:


A 1-factorisation of a graph  $G$  is called perfect if the union of any two of its 1-factors yields a Hamilton cycle of  $G$ .

Every 1-factorisation of  $K_4$  is perfect.


Every 1-factorisation of  $K_6$  is perfect.

But not every 1-factorisation of  $K_8$  is perfect.


Example: our first 1-factorisation of  $K_8$




$F_0$




$F_1$




$F_2$




$F_3$



$F_4$










$F_5$



$F_6$

Observe that  $F_0 \cup F_1$  yields a pair of 4-cycles  
 (actually, no  $F_i \cup F_j$  is Hamiltonian for this 1-factorisation).  
 Hence this 1-factorisation is not perfect.



Every pair of 1-factors yields an 8-cycle.  
Hence this 1-factorisation is perfect.

## Perfect 1-Factorisation Conjecture (Kotzig, 1963)

For each  $m \geq 2$ ,  $K_{2m}$  admits a perfect 1-factorisation.

### Theorem (Kotzig, 1963)

$GK_{2m}$  is perfect if and only if  $2m - 1$  is prime.

### Theorem (Anderson, 1973)

If  $m$  is prime, then the 1-factorisation  $GA_{2m}$  is perfect.

## Perfect 1-Factorisation Conjecture (Kotzig, 1963)

For each  $m \geq 2$ ,  $K_{2m}$  admits a perfect 1-factorisation.

### Theorem (Kotzig, 1963)

$GK_{2m}$  is perfect if and only if  $2m - 1$  is prime.

### Theorem (Anderson, 1973)

If  $m$  is prime, then the 1-factorisation  $GA_{2m}$  is perfect.

### Corollary

$K_{2m}$  has a perfect 1-factorisation if  $2m$  is one of 4, 6, 8, 10, 12, 14, 18, 20, 22, 24, 26, 30, 32, 34, 38, 42, 44, 46, 48, 54, 58, 60, 62, 68, 72, 74, 80, 82, 84, 86, 90, 94, 98, etc.

This leaves unsettled: 16, 28, 36, 40, 50, 52, 56, 64, 66, 70, 76, 78, 88, 92, 96, 100, etc.

## Other known P1Fs of $K_{2m}$ for small $m$

16: Kotzig and Anderson, 1974

There are 3155 nonisomorphic P1Fs of  $K_{16}$  (Gill and Wanless, 2020; Meszka, 2020)

28: Anderson, 1974

36: Seah and Stinson, 1988

40: Seah and Stinson, 1989

50: Ihrig, Seah and Stinson, 1987

52: Wolfe, 2009

This leaves unsettled: 56, 64, 66, 70, 76, 78, 88, 92, 96, 100, etc.

## Some other known P1Fs of $K_{2m}$

126, 170, 244, 344, 730, 1332, 1370, 1850, 2198, 3126, 6860, 12168, 29792 have been known since at least 1991. For references, see the survey by Seah in the *Bulletin of the ICA*, volume 1 (1991).

Several instances where  $2m = p^t + 1$  have been established. The most recent examples (by Wanless, 2005) include 530, 2810, 4490, 6890, 11450, 11882, 15626, 22202, 24390, 24650, 26570, 29930, etc. For more details see Wanless' website.

Also see the survey by Alex Rosa in *Mathematica Slovaca*, volume 69 (2019).

Still unsettled: 56, 64, 66, 70, 76, 78, 88, 92, 96, 100, etc.



## Starters:

A starter in  $\mathbb{Z}_{2t-1}$  consists of a set  $S$  of  $t - 1$  disjoint unordered pairs  $\{x_i, y_i\} \subset \{0, 1, \dots, 2t - 2\}$  such that for each  $d \in \{1, 2, \dots, 2t - 2\}$ , one of the  $t - 1$  pairs  $\{x_i, y_i\}$  satisfies either  $x_i - y_i \equiv d \pmod{2t - 1}$  or  $y_i - x_i \equiv d \pmod{2t - 1}$ .

## Example:

For  $t = 7$  consider the set  $S$  with these pairs:

$\{0, 1\}$  produces  $d$  values of 1 and 12 (mod 13)

$\{4, 6\}$  produces  $d$  values of 2 and 11 (mod 13)

$\{9, 12\}$  produces  $d$  values of 3 and 10 (mod 13)

$\{7, 11\}$  produces  $d$  values of 4 and 9 (mod 13)

$\{5, 10\}$  produces  $d$  values of 5 and 8 (mod 13)


$\{2, 8\}$  produces  $d$  values of 6 and 7 (mod 13)

## Starters:

Observe that a starter in  $\mathbb{Z}_{2t-1}$  yields a near 1-factorisation of  $K_{2t-1}$ .

## Example:

$$t = 7$$




## Starters:

Observe that a starter in  $\mathbb{Z}_{2t-1}$  yields a near 1-factorisation of  $K_{2t-1}$  and also a 1-factorisation of  $K_{2t}$ .

## Example:

$$t = 7$$



## Even Starters:

An even starter in  $\mathbb{Z}_{2t-2}$  consists of a set  $E$  of  $t - 2$  disjoint unordered pairs  $\{x_i, y_i\} \subset \{0, 1, \dots, 2t - 3\}$  such that for each  $d \in \{1, 2, \dots, 2t - 3\} \setminus \{t - 1\}$ , one of the  $t - 2$  pairs  $\{x_i, y_i\}$  satisfies either  $x_i - y_i \equiv d \pmod{2t - 2}$  or  $y_i - x_i \equiv d \pmod{2t - 2}$ .

## Example:

For  $t = 8$  consider the set  $E$  with these pairs:

$\{10, 11\}$  produces  $d$  values of 1 and 13 (mod 14)

$\{6, 8\}$  produces  $d$  values of 2 and 12 (mod 14)

$\{2, 5\}$  produces  $d$  values of 3 and 11 (mod 14)

$\{3, 7\}$  produces  $d$  values of 4 and 10 (mod 14)

$\{4, 13\}$  produces  $d$  values of 5 and 9 (mod 14)


$\{1, 9\}$  produces  $d$  values of 6 and 8 (mod 14)

## Even Starters:

Observe that an even starter in  $\mathbb{Z}_{2t-2}$  yields a 1-factorisation of  $K_{2t}$

## Example:

$$t = 8$$




## Even Starters:

Observe that an even starter in  $\mathbb{Z}_{2t-2}$  yields a 1-factorisation of  $K_{2t}$

## Example:

$$t = 8$$




## Even Starters:

Observe that an even starter in  $\mathbb{Z}_{2t-2}$  yields a 1-factorisation of  $K_{2t}$  when combined with this 1-factor.

## Example:

$$t = 8$$



## Wolfe's Approach for $K_{4m}$

Begin by finding a pair of starters in  $\mathbb{Z}_{2m-1}$ .

Merge them to build an even starter in  $\mathbb{Z}_{4m-2}$ .

Each of the  $m - 1$  pairs of each starter is given a high/low designation as part of the construction.

So in fact  $2^{m-1}$  even starters can be built from each pair of starters.

Use the even starter to build a 1-factorisation for  $K_{4m}$ .

Test the 1-factorisation for perfection.

Do this many times.



Wolfe found a P1F for  $K_{52}$ , published in 2009

He tested 7.494 billion pairs of starters in  $\mathbb{Z}_{25}$ .

Each pair produced  $2^{12} = 4096$  even starters in  $\mathbb{Z}_{50}$ .

To find one that yielded a P1F of  $K_{52}$  took 10,000 hours of computing time (i.e., about 166 days) on a cluster.

Real time was 5 days.

It had been about 20 years since the previous smallest open case of the Perfect 1-Factorisation Conjecture was settled.

## A P1F for $K_{56}$ , published in 2019

Wolfe's approach was used on a cluster with one director task and 1023 worker tasks running in parallel.

Each worker built pairs of starters in  $\mathbb{Z}_{27}$ , merged them in  $2^{13} = 8192$  ways, and tested the resulting 1-factorisations.

The worker that found a P1F had compared 7,730,443 pairs of starters in a real time span of 33 days 6 hours.

The other 1022 workers were terminated after 43 days 9 hours.

Estimated total number of pairs of starters: 10.3 billion

Total computing time for workers: 1,064,700 hours  
(i.e., a bit more than 121 years)

## A P1F for $K_{56}$ , published in 2019

The even starter of  $\mathbb{Z}_{54}$  shown below yields a P1F for  $K_{56}$ .

$\{36, 17\}, \{44, 12\}, \{39, 45\}, \{18, 35\}, \{8, 50\}, \{23, 15\},$   
 $\{42, 32\}, \{5, 46\}, \{19, 49\}, \{22, 37\}, \{10, 6\}, \{33, 30\}, \{3, 41\},$   
 $\{14, 21\}, \{48, 43\}, \{16, 52\}, \{25, 34\}, \{7, 38\}, \{11, 31\}, \{4, 2\},$   
 $\{29, 28\}, \{1, 27\}, \{0, 40\}, \{13, 24\}, \{51, 26\}, \{53, 20\}$

The smallest open case of the Perfect 1-Factorisation Conjecture is now  $K_{64}$ .

Thank you.

Acknowledgements:



Special thanks to the Centre for Health Informatics and Analytics.